Your thesaurus codes are: 07 (07.09.1; 03.01.1; 16.01.1; 04.01.1)

ASTRONOMY AND ASTROPHYSICS 17.7.2006

# First DENIS I-band extragalactic catalog \*

Vauglin I.,<sup>1</sup> Paturel G.,<sup>1</sup> Borsenberger J.,<sup>2</sup> Fouqué P.,<sup>3,4</sup> Epchtein N.,<sup>5,3</sup> Kimeswenger S.,<sup>6</sup> Tiphène D.<sup>3</sup> Lanoix P.,<sup>1</sup> Courtois H.,<sup>1</sup>

<sup>1</sup> CRAL-Observatoire de Lyon, F69561 Saint-Genis Laval CEDEX, FRANCE

<sup>2</sup> Institut d'Astrophysique de Paris
98 bis boulevard Arago, F75014 Paris, FRANCE

<sup>3</sup> Observatoire de Paris-Meudon F92195 Meudon Principal CEDEX, FRANCE

<sup>4</sup> European Southern Observatory La Silla, La Serena, CHILE

<sup>5</sup> Observatoire de Nice Departement Fresnel, BP 4429, F06304 Nice CEDEX, FRANCE

<sup>6</sup> Institut fur Astronomie Technikerstrasse 25, A6020 Innsbruck, AUSTRIA

Submitted August, 1998; Accepted

Abstract. This paper presents the first I-band photometric catalog of the brightest galaxies extracted from the Deep Near Infrared Survey of the Southern Sky (DENIS) An automatic galaxy recognition program has been developed to build this provisional catalog. The method is based on a discriminating analysis. The most discriminant parameter to separate galaxies from stars is proved to be the peak intensity of an object divided by its array. Its efficiency is better than 99%. The nominal accuracy for galaxy coordinates calculated with the Guide Star Catalog is about 6 arcseconds. The cross-identification with galaxies available in the Lyon-Meudon Extragalactic DAtabase (LEDA) allows a calibraton of the I-band photometry with the sample of Mathewson et Al. Thus, the catalog contains total I-band magnitude, isophotal diameter, axis ratio, position angle and a rough estimate of the morphological type code for 20260 galaxies. The internal completeness of this catalog reaches magnitude  $I_{lim} = 14.5$ , with a photometric accuracy of ~ 0.18m. 25% of the Southern sky has been processed in this study.

This quick look analysis allows us to start a radio and spectrographic follow-up long before the end of the survey. Key words: galaxies – catalog – photometry

## 1. Introduction

In Paturel et al. (1996), we presented our program of collecting the main astrophysical parameters for the principal galaxies. The first target was limited to adding information on galaxies already known in the LEDA database.

The work is now more ambitious because we are aiming at detecting new galaxies from the *Deep Near Infrared Survey of the Southern Sky* (hereafter DENIS). DE-NIS is a program to survey the entire southern sky in three wavelength bands (Gunn-i:  $0.80\mu m$ , J:  $1.25\mu m$  and Ks:  $2.15\mu m$ ) with limiting magnitudes of 18.5, 16.5 and 14.0 respectively. The observations are performed with the ESO 1m-telescope at La Silla (Chile) with a dedicated camera. The survey observations with three channels started in routine mode in December 1995. A detailed description of DENIS is given in Epchtein (1998) and in Garzón et al. (1997).

The systematic detection, extraction and cataloging of DENIS extragalactic sources are of significant interest for studies requiring large and homogeneous samples

Send offprint requests to: G. Paturel

<sup>\*</sup> Based on observations collected at the European Southern Observatory, La Silla, Chile.

such as the kinematics of the local universe, the distance scale, cosmology etc... The I-band is the most suitable both for the detection of extended objects and for the star/galaxy separation, except in the galactic plane. Thus, in the present study only the I-band images are considered.

The transfer of DENIS images from Paris to Lyon is explained in section 2 and the extraction of objects from these images in section 3. In sections 4 to 5 we describe astrometry and automatic galaxy recognition and analysis. Then, in sections 6 to 8, we explain how galaxies are crossidentified with LEDA galaxies leading to the comparison of astrophysical parameters with those from Mathewson et al. and from LEDA. Finally, in section 9 we describe the provisional I-band DENIS catalog.

It is important to note that a deeper catalog will be made at the end of the survey. Hence, the present catalog is a preliminary catalog of bright galaxiesdetected by DENIS, and used to start a radio and spectrographic follow-up long before the end of the survey. The present measurements cover one year of observation.

#### 2. Obtaining I-band CCD frames from DENIS

# 2.1. Characteristics of the DENIS survey

The I-band of the survey is the *Gunn-i* band at  $0.80\mu m$ . The CCD camera is a Tektronix  $1024 \times 1024$  pixel array cooled down to liquid nitrogen temperature. Each frame  $(768 \times 768 \text{ pixels})$  represents 12 square arcminutes with a pixel size of 1 arcsecond. The integration time is 9 seconds. The read-out noise is about  $7e^{-}$ . The observing strategy consists in scanning at a constant right ascension on strips of 30 degrees in declination (180 frames per strip) taken in three zones "Equatorial" from  $\delta = +2 \deg$  to  $-28 \deg$ , "Intermediate" from  $\delta = -28 \deg$  to  $-58 \deg$  and "Polar" from  $\delta = -58 \deg$  to  $-88 \deg$ . The overlap between adjacent frames is 1 arcminute on each side (i.e. 2 arcminute in each direction). This strategy aims at covering a wide range of airmasses. Each strip starts and ends with photometric and astrometric calibrations. At the end of some nights a flat fielding is performed directly on the sky during sunrise. The data is archived on DAT cartridges which are send each week to the Paris Data Analysis Center (PDAC) at the Institut d'Astrophysique de Paris for processing.

#### 2.2. Pipeline Lyon-PDAC

A systematic automatic processing of DENIS I-images began in Lyon in February 1996. Flat-fielding and de-biasing are made at PDAC on each genuine frame. For the Lyons processing each I-image is reduced by a factor 4 in size by rebinning pixels  $2 \times 2$ . Our effective pixel size is thus 2 arcseconds. An example of an I-frame is given in Figure 1.

The histogram of pixel intensities is used to calculate the sky background intensity  $f_{bg}$ . The sky-background level is taken at the maximum intensity of the histogram



**Fig. 1.** A typical image from the DENIS Survey after the  $2 \times 2$  rebinning. The frame is  $12' \times 12'$ . North is on the bottom side, East is on the left side. Among the four galaxies clearly visible on the top, one (the third from the left) is a new one. This shows that relatively bright galaxies can be discovered.

(i.e. the mode) and the standard deviation  $\sigma$  is calculated by symmetrizing the low intensity part of the histogram with respect to the mode. The sources are concentrated in the high intensity part of the histogram.

A threshold is then applied (the threshold level is  $f_{bg} + 2.0\sigma$ ). Using this procedure (averaging and thresholding) allows a compression factor of 20 to 30, depending on the image contents. All images of a given I-strip are thus compressed, tar'd and automatically transferred to Lyon via ftp. A full strip is stored in 10 to 13 Mbytes. The galaxy extraction is made at Lyon using the program described in the following section.

## 3. Extraction of astronomical objects

All sources (stars, galaxies, defects etc...) are extracted using the same algorithm as described in Paturel et al. (1996, section 3.1), except that no attempt was made to share interacting objects which are simply flagged after visual inspection (section 5). The reason is that we are interested first of all in well defined objects. At the end of this stage, we obtain for each frame a collection of matrixes (see an example in Figure 2). Matrixes smaller than 17 pixels are rejected. They correspond, to the mean, of objects of 6 arcseconds in diameter.

Astrophysical parameters are extracted for each matrix according to Paturel et al. (1996, section 3.2). These parameters are the following:



Fig. 2. Example of a matrix for a small galaxy. One pixel is  $2'' \times 2''$ . The edge is outside the matrix.

- Weighted mean position of the center  $x_m$  and  $y_m$  (in pixels), where the weight is the pixel intensity  $f_{ij}$
- major axis logD (D in 0.1arcmin.) at faint isophotal level (external diameter).
- axis ratio log R (log of major to minor axis)
- position angle of the major axis  $\beta$  (in degrees, counted from North towards East)
- magnitude  $I = -2.5 \log(\sum_{i=1}^{n} f_{ij} f_{bg}) + cst$  (in arbitrary units), n is the number of pixels with intensity  $f_{ij}$  larger than the threshold.

We have now to perform astrometry (conversion of pixels positions to right ascension and declination) and then recognition of "galaxies", "stars", and "unknown objects".

# 4. Astrometry

The center of the frame is taken from the header of the FITS file. From the Guide Star Catalog (GSC) and from the LEDA database we extract all objects (stars or galaxies) known in the corresponding square. A crossidentification between matrixes and stars is made exactly as described in Paturel et al. (1996, sections 3.3 and 3.4). Galaxies are also used but only when they have accurate coordinates (i.e. typically better than 10 arcsecond). A 6th order polynomial fit converts (x,y) positions on the frames to Right Ascension and Declination. The number of GSC stars varies from one frame to another. A histogram of number of GSC stars per frame is given in Figure 3. If this number is smaller than 7 or if the standard deviation of the polynomial fit is greater than 4 arcseconds, the solution is rejected and we adopt the 'header' solution calculated from the coordinates of the center and the pixel size as given in the header. If the GSC-solution seems acceptable but differs from the header-solution by more than 30 arcseconds, the header solution is preferred and coordinates are flagged to recall that they may be inaccurate.

In Figure 4, we show the differences between the GSCsolution and the FITS header-solution. Most of them are in good agreement within 15 arcseconds. Note that more



Fig. 3. Histogram of the number of GSC stars by frame.

recent measurements have been astrometrically calibrated to better than 1 arcsecond by cross-identifying with the PMM database.



**Fig. 4.** Histogram of the differences between the two astrometric solutions (GSC- and direct-solution).

Galaxy coordinates will be compared directly with coordinates of LEDA galaxies.

# 5. Automatic star/galaxy separation

#### 5.1. Test sample

A sample of 1146 objects has been visually classified into three classes: stars, galaxies and unknown objects. The distribution in each class is the following:

Class	Number of objects	percentage
Stars	500	43.6%
Galaxies	523	45.6%
Unknown	123	10.7%
Total	1146	100%

This sample will be used as a test sample (or training sample) in a discriminant analysis method (DA) for galaxy recognition.

#### 5.2. Discriminant analysis

The DA is a common method for automatic recognition. The test sample being shared in  $n_c$  classes  $\{I_k\}$  $(k = 1, n_c)$ , the purpose of DA is to find the principal factorial axis on which each class is as concentrated as possible and as distinct as possible from the others. This is achieved by maximizing the inertia between classes and by minimizing the inertia within each class. Inertia is calculated from the set of parameters attached to each object. We will note  $p_{ij}$  the *j*-th parameter of object *i*.

The mathematical result (see Diday et al., 1982) is that the factorial axes are the eigenvectors of the matrix  $\mathcal{T}^{-1}.\mathcal{B}$ , where  $\mathcal{T}$  is the total covariance matrix and  $\mathcal{B}$  is the inter-class covariance matrix.

Note that the matrix  $\mathcal{T}^{-1}.\mathcal{B}$  is not symmetrical and that the total covariance matrix is the sum of the intraclass covariance matrix  $\mathcal{W}$  (covariance Within class) and of inter-class covariance matrix  $\mathcal{B}$  (covariance Between class). This is called the Huyghens decomposition.

$$\mathcal{T} = \mathcal{W} + \mathcal{B} \tag{1}$$

The elements of  $\mathcal{B}$  are:

$$b_{jj'} = \sum_{k=1}^{n_c} \frac{N_k}{N} (\bar{p}_j^{\bar{k}} - \bar{p}_j) (\bar{p}_{j'}^{\bar{k}} - \bar{p}_{j'})$$
(2)

where  $N_k$  is the number of objects in the class  $I_k$  (k=1 to  $n_c$ ), where the mean parameter j for the whole sample is:

$$\bar{p_j} = \frac{1}{N} \sum_i p_{ij} \tag{3}$$

(N is the number of objects of the whole sample), and where the mean parameter j within the class  $\{I_k\}$  is:

$$\bar{p_j^k} = \frac{1}{N_k} \sum_{i \in I_k} p_{ij} \tag{4}$$

The elements of the total covariance matrix  $\mathcal{T}$  are:

$$t_{jj'} = \sum_{k} \frac{1}{N} \sum_{i \in I_k} (p_{ij} - \bar{p_j}) (p_{ij'} - \bar{p_{j'}})$$
(5)

Now, we have to choose the set of parameters attached to each object.

#### 5.3. Choice of discriminant parameters

Any discriminant method requires a good choice of discriminant parameters which are used for the definition of the metric. These parameters are not necessarily independent but they must cover all features which seem relevant for a reliable discrimination of astronomical objects. For galaxy recognition we tested 7 parameters.

- 1. Peak intensity per area unit, this is Peak intensity divided by the surface of the considered object.
- 2. Mean surface brightness, total flux divided by area
- 3. Peak intensity,
- 4. Axis ratio, ratio of the major to the minor axis
- 5. Relative area, ratio of number of pixels of the object and of the matrix.
- 6. Elongation of the matrix
- 7. Presence of diffraction cross

The DA method is applied on half the sample (i.e. 573 objects) and tested on the other half using only one parameter at a time (in this case the factorial axis is defined by the parameter itself). The percentage of good results is given below for each one, individually.

Parameter	stars	galaxies	mean
Peak over area	100.0%	98.9%	99.4%
Mean SB	98.7%	99.6%	99.2%
Peak intensity	96.2%	98.5%	97.4%
Axis ratio	83.7%	54.8%	69.2%
Relative surface	77.0%	58.5%	67.7%
Elongation of matrix	88.3%	40.8%	64.5%
Diffraction Cross	68.2%	57.0%	62.6%

The conclusion of this test is that the most relevant information about the nature of an object is contained in the pixel intensity, not in the shape of the object. Stars have a very high central intensity, galaxies do not. Moreover, stars are concentrated, galaxies are not. This explains why "Mean SB" and "Peak over area" give such an impressive recognition rate. Finally, only the first four parameters have been used. The axis ratio is kept because it becomes relevant for faint objects despite that its rate is relatively low.

### 5.4. Result

The DA method is applied with the four parameters described above and three classes "Galaxies", "Stars" and "unknown objects". Using the test sample, each object is projected onto the first factorial axis. Figure 5 shows the projection onto the first factorial axis of "Galaxies" and "Stars" classes. Similar plots exists for "Stars" and "unknown objects" classes and for "Galaxies" and "unknown objects" classes. All "unknown objects" have been eliminated in the next part of this study.

One can see that there is an overlapping region where "Galaxies" and "Stars" are mixed. The limits of this zone can be tuned in such a way that one can accept a given percentage of misclassification. We choose 0% chance of classifying a star as a galaxy and 5% chance of classifying a galaxy as a star. Indeed, it is important to avoid the contamination of the catalog by stars while it is not as important to miss a galaxy (which is uncertain anyway). These limits are drawn on Figure 5 where it is visible that no star enter the galaxy-domain, while 5% of galaxies enter the star-domain. Objects between these two limits will be classified as undefined.



Fig. 5. Definition of acceptation zones along the first factorial axis. The left-hand zone defines "Galaxies", the righthand one defines "Stars", and the intermediate one defines "undefined objects".

## 5.5. Visual control

The final step of this treatment consists in checking visually all frames recognized as galaxies. This tedious part allows us to reject artefacts (1148 rejections after the inspection of 54073 images) like those produced by star halos truncated by the edge of the frame. Such truncated halos look like elongated, low-surface brightness object, easily accepted as galaxies.

- As a result, a code is given to describe three features:
- "multiple", if several objects are present in the matrix
- "truncated", if the galaxy is truncated by the edge of the array.
- "peculiar", if the galaxy looks strange for any reason

So, each galaxy of the catalog has been inspected visually. This will prevent us from gross misidentification. Now, the galaxies have to be cross-identified with known galaxies.

## 6. Cross-identification

To make a correct cross-identification we need coordinates, calibrated magnitudes and diameters, axis ratios and position angles. We will use the first calibration obtained from the preliminary cross-identification made for astrometric purposes (section 4). This calibration will be refined in next section.

Because of frame overlap along the strip, many objects are measured twice (or even three or four times with adjacent strips). A first cross-identification is done for these galaxies measured several times. This will be called the "auto-cross-identification". Then, the cross-identification with LEDA galaxies may start.

# 6.1. Auto-cross-identification

The "auto-cross-identification" is performed using a hierarchical method in which we merge step by step the closest objects. The definition of the distance of two objects i and j is the following:

$$d_{ij} = \frac{1}{n_p} \sum_{k=1}^{n_p} \frac{|p_{ik} - p_{jk}|}{2\sigma_k}$$
(6)

where,  $n_p$  is the number of parameters (coordinates, diameters magnitudes...) for a given object,  $p_{ik}$  is the kth parameter of object i,  $\sigma_k$  is the standard error of the k - th parameter. When two objects are merged they are replaced by a single one, whose parameters are the means of both. The final result does not depend on the order the original file is read. Note that, special care must be taken for periodic parameters, Right ascension and position angle (e.g., p.a. = 0 deg is identical to p.a. = 180 deg; thus, the mean of p.a. = 3 deg and p.a. = 177 deg is p.a. = 0 deg, not p.a. = 90 deg).

The adopted  $\sigma_k$  are given in the following table:

Parameter	$\sigma_k$
$\alpha_{2000}$	30//
$\delta_{2000}$	30//
$\log D$	0.08
$\log R$	0.08
$I_T$	0.3 magnitude
$\beta$	$5/\log R$ degree

The standard error of the position angle is taken as a function of logR because its meaning vanishes for face-on galaxies.

Objects are merged for  $d_{ij} < d_{limit}$ ,  $d_{limit}$  beeing chosen from the distribution of all distances (Fig.6). By its definition,  $d_{ij}$  has the meaning of a Student's t-test devided by 2 (it is thus dimensionless). A 1 $\sigma$  criterion corresponds roughly to  $d_{limit} = 0.5$ . However, the value of  $\sigma$  attached to each parameter is somewhat arbitrary, so is the definition of  $d_{limit}$ . We adopted  $d_{limit} = 0.55$ . This choice is guided by the minimum observed in the histogram of  $d_{ij}$  (Fig.6).

During "auto-cross-identification" a provisional DE-NIS number (called RED) is given to each entry of the

z  $000^{+}_{000}$   $000^{+}_{000}$   $000^{+}_{000}$   $000^{+}_{000}$  0.5 1 1.5 distance for auto-cross-identification

**Fig. 6.** Distribution of "distances"  $d_{ij}$  between two DENIS objects (*i* and *j*). This graph is used to chose  $d_{limit}$  in the auto-cross-identification phase (see text).

catalog. When a galaxy appears several times in the catalog, each original set of measurements is identified with the same RED number. Each entry will be cross-identified independently with LEDA galaxies, this will allow us to check the reliability of the cross-identification.

#### 6.2. Cross-identification with LEDA galaxies

From the previous step we get an intermediate catalog in which each galaxy has a provisional number and all its astrophysical parameters ( $\alpha_{2000}$ ,  $\delta_{2000}$ ,  $log D_I$ ,  $log R_I$ ,  $I_T$ and  $\beta$ ). Each entry of this catalog must now be crossidentified with LEDA galaxies in order to identify those already known.

LEDA galaxies have similar parameters  $\alpha_{2000}$ ,  $\delta_{2000}$ ,  $logD_{25}$ ,  $logR_{25}$ ,  $B_T$  and  $\beta$ , but diameters and magnitudes are defined in the photometric B-band. The crossidentification is done by calculating the distance (in the mathematical sense, as defined by equation 6) between a DENIS and a LEDA galaxy. The coincidence is accepted if the distance is smaller than  $d_{limit}$ . From a histogram of all distances between LEDA and DENIS measurements (Fig.7), we adopted  $d_{limit} = 1.0$  which corresponds to the first minimum of  $d_{ij}$ -histogram (a pure Student's t-test would have given  $d_{limit} = 1.29$  at a 0.01 probability level). This limit is voluntarily conservative (i.e., small) because we prefer to miss a cross-identification than to merge two distinct galaxies.

Because a given galaxy is cross-identified each time it appears in the catalog, 15945 were cross-identified several times with their original parameters. We reject 1881 galaxies identified with more than one LEDA galaxy. The final catalog contains now 36247 galaxies.

Fig. 7. Distribution of "distances"  $d_{ij}$  between DENIS and LEDA galaxies (*i* and *j*). This graph is used to chose  $d_{limit}$  in the cross-identification phase (see text).

1.5

## 6.3. Last cleaning

Many objects were kept in the catalog despite the fact that they were labelled "undefined objects" by the DA. They were kept because a known galaxy was very close. In the present release we removed these objects which nature may be questionable without further inspection. Indeed, some very faint galaxies in LEDA have only a few parameters, so that the cross-identification is based mainly on coordinates. In some crowded field (clusters of galaxies or low galactic latitude) the accuracy of coordinates does not allow an identification secure enough. 10696 such objects were then rejected, leaving 25551 galaxies.

Finally, we rejected galaxies with uncertain coordinates so that only coordinates based on the GSC reference are used. So, 5291 galaxies are rejected in the present version, leading to the final catalog of 20260 galaxies. These drastic rejections aim at maintaining a high quality level for this first catalog. In order to judge the quality more quantitatively we will now compare with other sources of data.

## 7. Comparison with Mathewson et al. samples

## 7.1. Magnitudes

Magnitudes are calibrated by comparison with the measurements in I-band photometry made by Mathewson et al. (1992, 1996) which gives access to 2441 galaxies. This comparison allows us to correct for a possible variation of the zero-point. This variation has been explained by seasonal variation of the mean temperature of the camera <sup>1</sup>. Figure 8 shows such a variation described by the parameter C:



<sup>&</sup>lt;sup>1</sup> the camera is now equiped with a regulated cooling system.

Night number	С
1850 - 2150	+0.05
2151 - 2350	-0.08
2351 - 2600	+0.03
2600 - 2800	-0.05

Note that the "night number" is a logical number, not a real night number. Each time the system is initialized the night number is incremented. This explains that after one year of running survey there are 2500 logical nights.



Fig. 8. Zero-point variation obtained by comparison with Mathewson's I-band photometry.

An airmass correction is adopted using a typical value  $\Delta I/\Delta sec\zeta = 0.04 \pm 0.02$ . A check is made to control that there is no airmass residual. The residual is smaller than 0.01 magnitude. The DENIS I-magnitude is then:

$$I(DENIS) = -2.5log(\sum_{i=1}^{n} f_{ij} - f_{bg}) + 24.01 - 0.04sec\zeta + C$$
(7)

The zero-point distribution of I(Mathewson) - I(DENIS) is Gaussian (Fig.9) with a standard deviation of 0.2 magnitude. If we assume that the error is identical for both systems the mean error on DENIS extragalactic I-band magnitudes would be about 0.14 magnitude. Because the uncertainty on Mathewson et al. data is probably smaller, the uncertainty on our I-band magnitudes is about ~ 0.18 magnitude.

In Fig.10, the comparison between I(Mathewson) and I(DENIS) is shown for galaxies with secure identification and being neither "multiple" nor "truncated". The direct regression is:

$$I(Math.) = 1.05 \pm 0.02I(DENIS) - 0.54 \pm 0.24$$
(8)



Fig. 9. Zero-point distribution after a tiny seasonal correction

with the following standard deviation, correlation coefficient and number of objects:  $\sigma = 0.21$ ,  $\rho = 0.977 \pm 0.004$ , n = 163 after 11 rejections at  $3\sigma$ .

Stricto sensu, the slope is not significantly different from 1, and the zero-point is not significantly different from 0. So, we are keeping the conservative solution:  $I(Math.) \equiv I(DENIS)$ . Among the 10 rejected galaxies, 8 can be explained by localized poor photometric conditions (because they correspond to a loss of flux). The measurements of corresponding nights will be counted with half weight. Two nights were rejected (night 2475 and 2159) because they give rejections in the comparison of different photometric parameters.

# 7.2. Diameters ans Axis Ratios

Diameters and axis ratios are also compared with those of Mathewson et al. samples. These comparisons are given in Fig.11 and Fig.12, respectively. The direct regression are:

$$logD(Math.) = 0.96 \pm 0.04 logD(DENIS) + 0.04 \pm 0.04$$
(9)

with  $\sigma = 0.10$ ,  $\rho = 0.88 \pm 0.02$ , n = 170 after 4 rejections at  $3\sigma$ .

For the axis ratio, it is better to force the intercept to be zero as generally admitted (see de Vaucouleurs et al., 1976). This avoids to have negative axis ratio after the application of the regression. The result is thus:

$$logR(Math.) = 0.94 \pm 0.04 logR(DENIS)$$
(10)

with  $\sigma = 0.10$ ,  $\rho = 0.85 \pm 0.02$ , n = 172 after 2 rejections at  $3\sigma$ .

None of these regressions is significantly different from the absolute identity. So, we will keep:  $logD(Math.) \equiv$ logD(DENIS) and  $logR(Math.) \equiv logR(DENIS)$ . The standard deviations are 0.10 for both logD and logR. Again, if we assume the same error on both systems the error on logD and logR is about 0.07.



Fig. 10. Comparison of extragalactic I-band magnitudes from Mathewson et al. and from DENIS



Fig. 11. Comparison of I-band isophotal diameters from Mathewson et al. and from DENIS.



Fig. 12. Comparison of I-band axis ratios from Mathewson et al. and from DENIS

# 8. Comparison with LEDA galaxies

#### 8.1. Equatorial coordinates

The coordinates are compared with coordinates given in LEDA. Only coordinates known as 'accurate' in LEDA (i.e. with a standard deviation less than 10 arcseconds) are used. The plot of  $\Delta\delta$  and  $\Delta\alpha.cos\delta$  in Figure 13 shows that there is no systematic distorsion ( $\Delta$  means (DENIS) - (LEDA)). The standard deviation is 6.5 arcseconds and 6.7 arcseconds for  $\alpha.cos\delta$  and  $\delta$ , respectively. Assuming an error of the same amplitude in LEDA and DENIS coordinates gives an uncertainty of  $6.6/\sqrt{2}$  arcseconds for DENIS right ascension and declination, i.e. an uncertainty of 6.6 arcseconds for the position of a galaxy.

In fact, the plot of  $\Delta \alpha \cos \delta$  vs.  $\Delta \delta$  exhibited two abnormal zones with a systematic shift of about 30 or 40 arcseconds. This problem appeared near the zenith. Thus, only objects with coordinates based on the GSC stars are kept in the present version. In the final catalog, the coordinates will be obtained through a full astrometric solution (mosaicing frames along each strip and with adjacent ones) and this problem will be solved without rejecting galaxies.

# 8.2. Position angle

Position angle is important for studies on the orientation of galaxies, but also for identification. However, for nearly face on galaxies, it becomes very uncertain. The comparison is made only for galaxies with logR(DENIS) > 0.5.



Fig. 13. Comparison of coordinates from LEDA and DENIS.

The result is given in (Fig.14). The direct regression gives:

$$\beta(LEDA) = 1.009 \pm 0.003\beta(DENIS) + 0.07 \pm 0.26 \quad (11)$$

with  $\sigma = 2.65$ ,  $\rho = 0.9987 \pm 0.0001$ , n = 408 after 21 rejections at  $3\sigma$ . Among the 21 rejections, only 2 correspond to real discrepancies.

The uncertainty on the measurement of  $\beta$  is 2.7 deg.. This excellent agreement of position angles validates the reliability of our cross-identification.

## 8.3. Photometric morphological type code

It is interesting to obtain at least a very crude estimate of the morphological type code. This is particularly important when we plan to start a HI follow-up for which it is compulsory to avoid elliptical galaxies. The log of the standard deviation of pixel intensities is significantly correlated with the morphological type code extracted from LEDA. The relation is almost linear. However, it appears that the solution depends on the size of the galaxy (all very small objects appear identical). The slope and the intercept are found linearly correlated with the log of the number of pixels. In the scale between -5 and 10 defined in the Second Reference Catalog of Bright Galaxies, the photometric morphological type code is calculated as:

$$T(DENIS) = A \log(\sigma(f_{ij})) + B$$
(12)

Fig. 14. Comparison of position angles from LEDA and DE-NIS.

Where  $\sigma(f_{ij})$  is the standard deviation of flux of all pixels of the matrix representing the galaxy :

$$\sigma(f_{ij})) = \sqrt{\frac{\sum_{i,j} (f_{ij} - \bar{f})^2}{(n-1)}}$$
(13)

with

$$A = \frac{1}{-0.0187 \log n - 0.0183} \tag{14}$$

$$B = \frac{1}{0.2774 \log n + 1.7846} \tag{15}$$

where n is the number of pixels.

The comparison of codes T(DENIS) and T(LEDA)is given in Figure 15. The correlation is clearly significant (the correlation coefficient is  $\rho = 0.69 \pm 0.01$ ) but the standard deviation is large ( $\sigma(T(DENIS)) = 2.5$ ). This allows us to classify galaxies into 'Early', 'Intermediate' and 'Late' types, with no finer subdivision. In the catalog, photometric morphological type smaller than -5 or greater than 10 will be set to -5 or 10, respectively.

## 9. The catalog

# 9.1. Description of the catalog

The catalog of galaxies which is described here is the first of a series which will be released during the progression of the survey. At the end of the survey a deeper and complete





Fig. 15. Comparison of morphological type codes from LEDA and from the I-band photometry.

catalog will be produced at PDAC. The present one, is necessarily limited to the first observations (one year).

The catalog contains 20260 galaxies. Among them 14518 are new galaxies while 5742 are galaxies already present in LEDA.

Each galaxy is numbered with a provisional internal number (RED, for Rapid Extraction from DENIS), and an identifier from the LEDA database (PGC/LEDA) <sup>2</sup>. Galaxies are also identified with an alternate name taken in the following catalogues: NGC, IC (Dreyer, 1988), UGC (Nilson, 1973), ESO (Lauberts, 1982), MCG (Vorontsov-Velyaminov et al., 1962-1974), CGCG (Zwicky et al., 1961-1968), IIZW, IIIZW, VIII from the catalog of compact and eruptive galaxies (Zwicky, 1971), IISZ (extension of the same list by Rodgers et al., 1978, FAIR (Fairall, 1977-1988), HICK (Hickson, 1989), KUG (Takase and Miyauchi-Isobe, 1984-1989), IRAS (IRAS), MK (Markarian, 1971-1977), UM (University of Michigan list, MacAlpine et al., 1981). The detailed references to these catalogs are given in Paturel et al. (1989). Two additional catalogs are included: Saito et al. (1990-1991) and Dressler (1980). The notations for corresponding galaxy names are SAIT (e.g. SAIT 69-1, for object 69 in list 1 of Saito et al.) and DRCG (e.g., DRCG 39-41, for galaxy 41 in cluster 39, the numbering of clusters is made according to the order in the original publication).

For each galaxy, the catalog gives the weighted mean parameters obtained as described in the previous section. Actual mean errors  $\sigma$  are calculated according to Paturel et al. (1996) using individual mean errors deduced in previous section. For I-band magnitudes the mean error is taken as a function of the magnitude itself. An esti-

mate made from Fig. 10 gives: $\sigma(I) = 0.05B_T - 0.51$ . The weights used for calculating the weighted means are the inverse square of actual mean errors. For nights suspected to be of poor photometric quality the actual mean error is divided by 3<sup>3</sup>. Further, when the quality of the matrix is uncertain ('peculiar', 'multiple' or 'truncated'), the weight is divided by 2. This correcting coefficient is deduced from the comparison of I magnitudes with poor quality objects.

The quality of each individual matrix is coded as follows: 'Normal' galaxies are coded as 1, 'Peculiar' galaxies as 10, 'Multiple' as 100 and 'Truncated' as 1000. This code is added for each measurement. So, the resulting code gives immediately the number of independent measurements (sum of digits) and the quality of each of them. For instance, the code 1002 means that there are one truncated matrix and two normal ones (i.e., 3 independent measurements). 'Truncated' means an overestimated magnitude, on the contrary, 'Multiple' means that the magnitude of the considered galaxy is underestimated.

The different columns are the following:

- Column 1:Provisional DENIS identifier. This identifier is just an internal number given for easy identification. This number will be replaced by an appropriate final DENIS number.
- Column 2:LEDA identifier. This number allows rapid retrieval in the LEDA database. This number is identical to the PGC number for LEDA number less than 73197.
- Column 3: Alternate name according to a hierarchy: NGC, IC, ESO, UGC, MCG (see text above).
- Column 4:Equatorial coordinates for epoch 2000, in hours, minutes, seconds and tenths for the Right Ascension and in degrees, arcminutes and arcseconds for the declination. The actual mean error on the position of the galaxy is given in the same column, next line in arcseconds.
- Column 5: Total I-band magnitude I(DENIS) with its mean error on the second line.
- Column 6: Isophotal diameter in log scale logD(DENIS), where D is in 0.1 arcminute following the notation of de Vaucouleurs et al. (RC1, RC2). The actual mean error is given on the second line.
- Column 7: Isophotal axis ratio in log scale logR(DENIS), where R is the ratio of the major to the minor axis. The actual mean error is given on the second line.
- Column 8: Position angle  $\beta(DENIS)$  in degrees, measured from North Eastwards. Its value lays between 0 and 180 degrees (excluding 180). For logR(DENIS) < 0.5 the meaning is poor. This is reflected by the actual mean error given on the second line.

<sup>&</sup>lt;sup>2</sup> connection:

http://www-obs.univ-lyon1.fr/leda or

telnet: lmc.univ-lyon1.fr login:leda

 $<sup>^3</sup>$   $\,$  These nights were detected by a  $3\sigma$  rejection rule, so one can admit that their mean error is three times the typical mean error.

- Column 9: Photometric morphological type code T(DENIS). The coding is given according to RC2 (i.e. T from -5 to 10 for Elliptical to Irregular galaxies). The mean error is given on the second line.
- Column 10: Quality code of the object (1000=truncated object, 100=multiple object, 10=peculiar object, 1=normal galaxy). The sum of digits gives the number of independent measurements.

A sample page is given at the end of the paper. The catalog is available in electronic form at PDAC and Lyons Observatory.

#### 9.2. Distribution on the sky

The distribution on the sky is given in equatorial coordinates for the epoch 2000 (Figure 16). The strips are clearly visible. The zone called 'Equatorial' ( $\delta$  between +2 deg and -28 deg) is not so well covered because observations are avoided in this zone when Moon is bright or when the wind is strong. Further, many frames are rejected in this zone at the end of the strip (near -28 deg) because of the shift in header-coordinates. No attempt is made to reach low galactic latitudes. This is done independently in J and K bands.

# 9.3. Completeness limit

If we assume a homogeneous distribution of galaxies ( $N \propto r^3$ ), the plot of the number of galaxies (in log scale, i.e. logN) brighter than a given magnitude limit  $I_{limit}$  versus this magnitude limit should be linear with a slope of 0.6. This plot (Figure 17) is quite linear up to  $I_{limit} = 14.5$  with a slope of  $0.62 \pm 0.02$ . The sense of this completeness limit must be explained. It means that in surveyed directions all galaxies brighter than 14.5 are detected. Because the sampling is made randomly and 1/4 of the survey is presented here, the completeness limit in apparent magnitude of this first DENIS catalog is  $I_{limit} = 14.5$ .

Acknowledgements. This work would have been impossible without the collaboration of Marthinet M.C., Petit C., Provost L., Gallet F., Garnier R., Rousseau J. They are fully associated with this work. The DENIS team is warmly thanked for making this work possible and in particular the Operations team at la Silla. We thank also G. Mamon for careful reading of the article.

The DENIS program is partly funded by the European Commission through *SCIENCE* and *Human Capital and Mobility* grants. It is also supported in France by INSU, the Education Ministry and CNRS, in Germany by the Land of Baden-Würtenberg, in Spain by DGICYT, in Italy by CNR, in Austria by the Fonds zur Förderung der Wissenschaft und Forschung, in Brazil by FAPESP.

#### References

Diday E., Lemaire J., Pouget J., Testu F., 1982, 'Eléments d'analyse de données', Eds. Dunod, Paris, ISBN 2-04-015430-2



Fig. 17. Completeness curve. The completeness curve is quite linear up to the completeness limit  $I_{limit} = 14.5$ . The slope  $(0.62 \pm 0.02, \text{ dashed line})$  is very nearly the one expected for a homogeneous distribution of galaxies (0.6).

Dressler A.; 1980, Astrophys. J. Suppl. Ser., 42, 565

- N. Epchtein, (ed.), 1998, 'The impact of near infrared surveys on galactic and extragalactic astronomy', Proc. 3rd. Euroconf. Kluwer ASSL series 230
- Epchtein, N. + 47 authors, 1997, The Messenger No 87, p.27
- Garzón F., Epchtein N., Omont A., Burton W.B., Persi P. (eds.), 1997, 'The impact of large-scale near-infrared sky surveys', ASSL series vol. 210, Kluwer Acad. Pub., Dordrecht
- Guide Star Catalog, 1989-1992, Space Telescope Science Institute (GSC)
- Lebart L., Fénelon J.P., 1975, 'Statistique et Informatique Appliquées', Eds. Bordas, Paris
- Mathewson D.S., Ford V.L., Buchhorn M., 1992, Astrophys. J. Suppl. Ser., 81, 413
- Mathewson D.S., Ford V.L., 1996, Astrophys. J. Suppl. Ser., 107, 97
- Paturel G., Fouqué P., Bottinelli L., Gouguenheim L., 1989, Astron. Astrophys. Suppl.Ser. 80,299
- Paturel G., Garnier R., Petit C., Martinet M.C., 1996, Astron. Astrophys. 311, 12
- Paturel G., Andernach H., Bottinelli L., Di Nella H., Durand N., Garnier R., Gouguenheim L., Lanoix P., Marthinet M.C., Petit C., Rousseau J., Theureau G., Vauglin I., 1997, Astronom. Astrophys. Suppl. Ser. 124, 109
- Saito M. et al.; 1990, PASJ, 42, 565
- Saito M. et al.; 1991, PASJ, 43, 449
- de Vaucouleurs G., de Vaucouleurs A., Corwin H.G. Jr., 1976, Second Reference Catalog of Bright Galaxies, University of Texas Press, Austin (RC2)
- de Vaucouleurs G., de Vaucouleurs, A., Corwin, H.G. Jr., Buta, R.J., Paturel, G., Fouqué, P., 1991, Third Reference Catalog of Bright Galaxies, Springer-Verlag (RC3)



Fig. 16. Distribution of the 20260 galaxies of the present DENIS catalog presented with a Flamsteed's equal area projection in equatorial (2000) coordinates.

This article was processed by the author using Springer-Verlag  ${\rm IAT}_{\rm E}{\rm X}$  A&A style file L-AA version 3.

TABLE 1: Sample page of the present catalog.

DENIS-RED (1)	LEDA/PGC (2)	Alte	rnate Name (3)	R.A. DEC.(2000) (4)	I (5)	logD (6)	logR (7)	p.a. (8)	T (9)	code (10)
000142	000087	DRCG	39- 41	J000109.8-503505	14.82	.718	. 306	42.9	0.	2
000149	143168			5 J000110.8-404542	.17 12.97	.061 .873	.057 .083	5.5 17.0	2. -4.	1
000150	143169			6 J000111.7-384507	.14 15.45	.070	.070	19.0 102.0	3. 6.	1
000153	124807			6 J000112.3-391354	.26 14.99	.648	.188	3.9 95.9	3. -1.	2
000159	143172			4 J000115.9-730851	.21 15.28	.052 .683	.080 .417	21.5 120.0	3. -2.	1
000160	000099	ESO	149- 12	6 J000117.5-530033	.36 13.54	.099 1.138	.099 .701	6.4 36.9	4. 4.	2
000165	143175			5 J000119.7-523551	.12 14.29	.050 .828	.054 .389	2.2 14.1	2. 10.	2
000168	143177			5 J000120.4-404721	.15 14.70	.067 .713	.051 .236	4.0 44.0	2. 1.	1
000170	143179			6 J000121.0-723533	.23 14.97	.070 .683	.070 .181	7.7 134.0	3. 0.	1
000174	130913			6 J000131.1-404912	.24 13.99	.070 .873	.070 .208	9.9 114.0	3. 10.	1
000182	124990			6 J000138.5-435949	.19 14.51	.070	.070	8.7 153.0	3. -2.	1
000183	124811			6	.22	.070	.070	90.9	3.	1
000105	142200			6 10000000 1 F1F740	.27	.070	.070	7.0	3.	-
000220	143200			6	.22	.070	.070	9.9	3.	I A
000253	143209			5000218.6+015033	.22	.683	.181	9.9	-4. 3.	1
000272	143215			J000233.8-111705 6	14.31 .21	.903 .070	.458 .070	78.0 4.2	3. 3.	1
000289	000205	UGC	5	J000305.7-015449 5	12.18 .07	1.248 .052	.354 .069	51.5 4.2	5. 2.	2
000292	143223			J000307.6-160643 6	14.64 .22	.723 .070	.153 .070	90.0 11.4	3. 3.	1
000293	143224			J000308.6-170300	14.55 22	.683	.125	83.0 13.6	-2.	1
000294	143225			J000309.0-195221	15.00	.703	.431	19.0	-5.	1
000296	000211			J000310.5-544457	.24	1.135	.167	4.4 58.1	-4.	1001
000299	143226			6 J000313.5-512806	.10 14.78	.071 .743	.078 .417	9.9 142.0	3. 0.	1
000303	143228			6 J000318.3-131618	.23 14.16	.070 .923	.070 .417	4.6 83.0	3. 10.	1
000307	143229			6 J000320.4-394823	.20 14.96	.070 .783	.070 .486	4.6 131.0	3. 3.	1
000309	000224	FAIR	627	6 J000321.3-500448	.24 13.47	.070 .933	.070 .347	3.9 110.2	3. -4.	2
000310	073217			5 J000321.4-543338	.12 12.81	.053 1.023	.109 .125	3.9 145.0	2. -2.	100
000311	143230			20 J000321.6-190604	.39 14.74	.210 .733	.210 .306	40.8 146.0	9. -2.	1
000313	143232			6 J000322.4-434615	.23 14.73	.070 .693	.070 .208	6.1 165.0	3. -2.	1
				6	. 23	.070	.070	8.7	3.	